arXiv:1201.4775v2 [math.RT] 8 Feb 2012

COMPUTATIONS FOR COXETER ARRANGEMENTS AND SOLOMON'S DESCENT ALGEBRA II: GROUPS OF RANK FIVE AND SIX

MARCUS BISHOP, J. MATTHEW DOUGLASS, GÖTZ PFEIFFER,
AND GERHARD RÖHRLE

Abstract. In recent papers we have refined a conjecture of Lehrer and Solomon expressing the character of a finite Coxeter group W acting on the $\mathfrak{p}^{\text{th}}$ graded component of its Orlik-Solomon algebra as a sum of characters induced from linear characters of centralizers of elements of W. Our refined conjecture relates the character above to a component of a decomposition of the regular character of W related to Solomon's descent algebra of W. The refined conjecture has been proved for symmetric and dihedral groups, as well as finite Coxeter groups of rank three and four.

In this paper, the second in a series of three dealing with groups of rank up to eight (and in particular, all exceptional Coxeter groups), we prove the conjecture for finite Coxeter groups of rank five and six, further developing the algorithmic tools described in the previous article. The techniques developed and implemented in this paper provide previously unknown decompositions of the regular and Orlik-Solomon characters of the groups considered.

1. Introduction

Let (W, S) be a finite Coxeter system and let T be the set of reflections in the complex reflection representation V of W. In previous articles [2–4] we proposed a conjecture relating the character ω of the Orlik-Solomon algebra of W to the regular character ρ of W. In this paper we prove the conjecture for the Coxeter groups of type B_5, B_6, D_5, D_6, and E_6. These computations, together with the remarks about reducible Coxeter groups following Theorem 2.3 of [2] and the proof of the conjecture for groups of type A in [3], prove the conjecture for all finite Coxeter groups of rank five or six. Our result is stated for these groups as the following theorem.

Theorem 1. *Suppose that W is a finite Coxeter group of rank five or six and that $\mathcal{R}$ is a set of conjugacy class representatives of W. Then for each $w \in \mathcal{R}$ there exists a linear character φ_w of $C_W(w)$ such that*

$$\rho = \sum_{w\in\mathcal{R}} \operatorname{Ind}_{C_W(w)}^{W} \varphi_w \quad \textit{and} \quad \omega = \epsilon \sum_{w\in\mathcal{R}} \operatorname{Ind}_{C_W(w)}^{W} (\alpha_w \varphi_w)$$

2010 *Mathematics Subject Classification.* 20F55, 20C15, 20C40, 52C35.

Key words and phrases. Coxeter group, descent algebra, Orlik-Solomon algebra.

where ϵ is the sign character of W and α_w is the composition of det *with restriction to the* 1*-eigenspace of* w *in* V.

Recall that the Orlik-Solomon algebra of W is the quotient of the exterior algebra with generators $\{e_t \mid t \in T\}$ modulo the ideal generated by elements of the form $\sum_{i=1}^{k} (-1)^i e_{t_1} e_{t_2} \cdots \widehat{e_{t_i}} \cdots e_{t_k}$ for all sets $\{t_1, t_2, \ldots, t_k\} \subseteq T$ of linearly dependent reflections. Here, we say that a set of reflections is linearly dependent if the forms defining their reflecting hyperplanes are linearly dependent in the dual of V. For $t \in T$ we denote the image of the generator e_t in $A(W)$ by a_t. Thus, an arbitrary element of $A(W)$ can be expressed as a linear combination of monomials $a_{t_1} a_{t_2} \cdots a_{t_k}$ with $t_1, t_2, \ldots, t_k \in T$. Because conjugating a reflection yields another reflection, $A(W)$ is a right $\mathbb{C}W$-module with $(a_{t_1} a_{t_2} \cdots a_{t_k}).w = a_{w^{-1}t_1 w} a_{w^{-1}t_2 w} \cdots a_{w^{-1}t_k w}$ for $w \in W$.

The strategy for proving Theorem 1 is to decompose the representations $\mathbb{C}W$ and $A(W)$ into direct sums and prove an analog of Theorem 1 for each summand. This method is somewhat stronger than directly proving Theorem 1 because it requires the solution to be compatible with the direct sum decompositions of $\mathbb{C}W$ and $A(W)$. However, it has the advantage that it splits the problem into many smaller problems and provides insight into how the representation of W on $A(W)$ and the right regular representation of W are related.

The decomposition of $\mathbb{C}W$ comes from the idempotents e_λ in the descent algebra $\Sigma(W)$ constructed in [1]. These idempotents are indexed by subsets of S up to conjugacy in W. Such an equivalence class is called a *shape* of W. In fact, [1] shows how to construct a quasi-idempotent e_L for any $L \subseteq S$. Then e_λ is the sum of the quasi-idempotents e_L where L runs over the subsets in the shape λ. It is also shown that $\{e_\lambda \mid \lambda \in \Lambda\}$ is a complete set of primitive orthogonal idempotents of $\Sigma(W)$, where Λ is the set of all shapes of W. Since $\Sigma(W)$ is a subalgebra of $\mathbb{C}W$, we conclude that $1 = \sum_{\lambda \in \Lambda} e_\lambda$ so that $\mathbb{C}W = \bigoplus_{\lambda \in \Lambda} \mathbb{C}We_\lambda$, as W-module. Denoting the character of $\mathbb{C}We_\lambda$ by ρ_λ we have $\rho = \sum_{\lambda \in \Lambda} \rho_\lambda$.

The decomposition of $A(W)$ comes from organizing its monomials according to their fixed point spaces as follows. The monomial $a_{t_1} a_{t_2} \cdots a_{t_k}$ determines a subspace of V, namely the intersection of the fixed point spaces of the reflections $t_1, t_2, \ldots, t_k$. If X is a subspace of V, then we denote the span of all monomials whose fixed point spaces are exactly X by A_X. Then taking A_λ to be the direct sum of the A_X for which X is the fixed point set of W_L for some $L \in \lambda$, we have a decomposition $A(W) = \bigoplus_{\lambda \in \Lambda} A_\lambda$. Denoting the character of A_λ by ω_λ we have $\omega = \sum_{\lambda \in \Lambda} \omega_\lambda$.

Finally, we organize the set of conjugacy class representatives $\mathcal{R}$ into subsets corresponding with shapes. This organization is defined using the notion of *cuspidal conjugacy classes* in the parabolic subgroups $W_L = \langle L \rangle$ of W. A conjugacy class in W_L is called *cuspidal* if the fixed point set in the reflection representation of W_L of any of its elements is trivial. Now if we choose a fixed representative $L(\lambda)$ of each shape λ and let $\mathcal{C}_{L(\lambda)}$ be a set of representatives of the cuspidal classes in

$W_{L(\lambda)}$, then by Theorem 3.2.12 of [5], $\mathcal{R} = \bigcup_{\lambda\in\Lambda} \mathcal{C}_{L(\lambda)}$ is a set of conjugacy class representatives of W.

Suppose $L \subseteq S$ and consider the homogeneous component of $A(W_L)$ of highest degree. This component is a W_L-submodule of $A(W_L)$, so it affords a character ω_L which we call the *top component character* of $A(W_L)$. On the other hand, W_L is also a Coxeter group and hence has its own quasi-idempotents as in [1], now denoted by e^L_K for $K \subseteq L$ to distinguish them from the quasi-idempotents e_K in $\mathbb{C}W$. (Notice that e^L_K is not the idempotent denoted by e^L_K in [1, §7].) Then $\mathbb{C}W_L e^L_L$ is a submodule of $\mathbb{C}W_L$, so it affords a character ρ_L of W_L which we call the *top component character* of $\mathbb{C}W_L$.

Now $A(W_L)$ and $\mathbb{C}W_L e_L$ are naturally subspaces of $A(W)$ and $\mathbb{C}W$ and it turns out that they are $N_W(W_L)$-stable by [3, Proposition 4.8]. Thus they afford characters $\widetilde{\rho_L}$ and $\widetilde{\omega_L}$ of $N_W(W_L)$, which are extensions of ρ_L and ω_L. Furthermore, if $L = L(\lambda)$, then by the same proposition, $\rho_\lambda = \mathrm{Ind}^W_{N_W(W_L)} \widetilde{\rho_L}$ and $\omega_\lambda = \mathrm{Ind}^W_{N_W(W_L)} \widetilde{\omega_L}$.

Note that by [9, Theorem 3.1], if w is cuspidal in W_L, then $C_W(w)$ is a subgroup of $N_W(W_L)$. Therefore, if we can find a linear character $\widetilde{\varphi}_w$ of $C_W(w)$ for each $w \in \mathcal{C}_L$ such that inducing the characters $\widetilde{\varphi}_w$ and $\alpha_w \epsilon \widetilde{\varphi}_w$ to $N_W(W_L)$ and summing over the cuspidal classes in W_L yields $\widetilde{\rho_L}$ and $\widetilde{\omega_L}$ respectively, then by transitivity of induction, the characters $\widetilde{\varphi}_w$ satisfy the conclusion of Theorem 1 with ρ, ω, and $\mathcal{R}$ replaced by ρ_λ, ω_λ, and $\mathcal{C}_{L(\lambda)}$, respectively, where λ is the shape containing L.

Repeating the preceding discussion for all shapes λ shows that Theorem 1 is a consequence of the following theorems. Theorem 2 is proved in Section 3 and Theorem 3 is proved in Section 4.

Theorem 2. *Suppose that (W, S) is a finite Coxeter system of rank five or six and that $\mathcal{C}$ is a set of representatives of the cuspidal conjugacy classes of W. Then for each $w \in \mathcal{C}$ there exists a linear character φ_w of $C_W(w)$ such that*

$$\rho_S = \sum_{w\in\mathcal{C}} \mathrm{Ind}^W_{C_W(w)} \varphi_w = \epsilon\omega_S.$$

Theorem 3. *Suppose that (W, S) is a finite Coxeter system of rank five or six, that L is a proper subset of S, and that $\mathcal{C}_L$ is a set of representatives of the cuspidal conjugacy classes of W_L. Then for each $w \in \mathcal{C}_L$ there is a linear character $\widetilde{\varphi}_w$ of $C_W(w)$ such that*

$$\widetilde{\rho_L} = \sum_{w\in\mathcal{C}_L} \mathrm{Ind}^{N_W(W_L)}_{C_W(w)} \widetilde{\varphi}_w = \alpha_L \epsilon \widetilde{\omega_L},$$

where α_L is the composition of det *with restriction to the subspace of fixed points of W_L in V.*

Notice that if Theorem 2 has been proved for a parabolic subgroup W_L of W, then Theorem 3 could be proved for the same pair W, W_L by showing that each character φ_w of $C_{W_L}(w)$ satisfying Theorem 2 extends to a character $\widetilde{\varphi}_w$ of $C_W(w)$ satisfying Theorem 3. This is the line of reasoning used in [3] and [4]. For the cases not covered in these two references, in this paper the character $\widetilde{\varphi}_w$ is constructed directly as a character of $C_W(w)$ and not as an extension of a character of $C_{W_L}(w)$. However, as noted in [4, Proposition 4.5], as w varies over $\mathcal{C}_L$, the restrictions $\widetilde{\varphi}_w|_{C_{W_L}(w)}$ satisfy the conclusion of Theorem 2 for the group W_L.

2. Implementation

As in [2], we have implemented the calculations for this article in the computer algebra system GAP in conjunction with the CHEVIE [6], ZigZag [10], and Gauss [7] packages. In addition to our comments about the implementation in [2] we make the following remarks. Let W be a finite Coxeter group and let F an extension of $\mathbb{Q}$ containing the entries of the matrices in the reflection representation V of W. The calculations described below can all be carried out in a vector space over F and hence are possible to implement in a computer algebra system. Although F appears to play no role in the calculations below, its role is implicit whenever vector spaces are used, since GAP requires the user to specify the coefficient field of a vector space.

2.1. **The Extension $\widetilde{\rho_L}$.** Throughout the following calculation, we fix a list $E = (w_1, w_2, \dots, w_{|W|})$ of the elements of W and represent elements of FW as coefficient vectors with respect to E. Thus the element $\sum_{i=1}^{|W|} \alpha_i w_i$ of FW is expressed as the vector $(\alpha_1, \alpha_2, \dots, \alpha_{|W|})$. Representing elements of FW as coefficient vectors proves to be much faster than working with GAP's ring-theoretic functionality, even at the expense of having to occasionally search the list E, which can be quite long.

To construct the characters $\widetilde{\rho_L}$ for $L \subseteq S$, the first step is to construct parabolic transversals, x_L, in $\mathbb{C}W$. By definition, x_L is the sum of the elements w in W satisfying $\ell(sw) > \ell(w)$ for all $s \in L$, where ℓ is the usual length function of W with respect to S. Then x_L can be calculated directly from the definition or by using the `ParabolicTransversal` function supplied by the ZigZag package. We calculate the elements x_L for all $L \subseteq S$ and store them as coefficient vectors.

The quasi-idempotents e_L are defined in [1] by means of the matrix $M = (m_{KJ})$ whose rows and columns are indexed by the subsets of S and whose (K, J)-entry is

$$m_{KJ} = \begin{cases} \left|\left\{x \in X_J \mid x^{-1}Jx \subseteq K\right\}\right| & \text{if } J \subseteq K \\ 0 & \text{otherwise} \end{cases}$$

for all $J, K \subseteq S$. Then putting $N = M^{-1} = (n_{JK})$ we have $e_L = \sum_{J \subseteq S} n_{KJ} x_J$. In terms of coefficient vectors, e_L can be calculated by multiplying the matrix N by the matrix of coefficient vectors of the elements x_L in the same order as the rows and columns of M. The matrix M can be calculated directly from the definition or by calling the method `Mu` supplied by the ZigZag package.

With e_L expressed as a coefficient vector v, we next calculate a basis of the $N_W(W_L)$-stable subspace FW_Le_L. Each element w of W_L acts on a coefficient vector by some permutation π_w, which can be determined by simply multiplying w by all the elements of W and recording the positions in E of the products. It follows that the vectors $v.\pi_w$ for all $w \in W$ span a subspace of $F^{|W|}$ isomorphic to $FW_L e_L$. Let B be a basis of this vector space.

To complete the calculation of the character $\widetilde{\rho_L}$, we choose conjugacy class representatives for $N_W(W_L)$ and for each representative w we compute the trace of the matrix of w with respect to the basis B.

2.2. The Extension $\widetilde{\omega_L}$.

To calculate the character $\widetilde{\omega_L}$ for $L \subseteq S$ we begin with the broken circuit basis of the algebra $A(W_L)$ described in [2]. This calculation depends only on the isomorphism type of W_L and is typically a very short calculation in comparison with the corresponding calculation for $A(W)$. However, care must be taken whenever elements of $N_W(W_L)$ act on $A(W_L)$, because in GAP the reflections of W_L in its reflection representation are typically indexed differently than the reflections of W in V.

Rather than working with reflections, it is equivalent and far simpler to work with positive roots, which in CHEVIE are stored as vectors in the `roots` component of a group record, the first half being the positive and the second half the negative roots. This means that whenever applying a reflection to a root results in a negative root, we need to replace the negative root with its positive counterpart. Now since CHEVIE implements the element w of W as a permutation σ_w of the roots, it follows that if t is the reflection defined by the root r, then the conjugate $w^{-1}tw$ is the reflection defined by $r.\sigma_w$. However, if w is an element of W and r is a root of W_L, then r must be replaced with its corresponding root of W using the `rootInclusion` component of the W_L record. Then the permutation σ_w can be applied directly. Now recording the monomial $a = a_{t_1} a_{t_2} \cdots a_{t_k}$ as the tuple $t_1, t_2, \ldots, t_k$ we can calculate the action of w on a by simply applying σ_w to each t_i and replacing any negative root with its positive counterpart. We use the method described in [2] to express arbitrary elements of $A(W_L)$ as coefficient vectors with respect to the broken circuit basis.

Finally, the set of $|L|$-tuples in the broken circuit basis of $A(W_L)$ form a basis B of the top component of $A(W_L)$. Then as in subsection 2.1, for each congacy class representative w for $N_W(W_L)$, we compute the trace of the matrix of w with respect to the basis B.

We also remark that in contrast with those in subsection 2.1, the matrices of the representation above are extremely sparse. While the matrices required for the calculation of any character $\widetilde{\omega_L}$ in this paper were not big enough to warrant using sparse matrices, the calculation of ω_S in Section 3 benefited greatly by using sparse matrices as implemented in the Gauss package for GAP [7].

3. PROOF OF THEOREM 2

In this section we exhibit the characters φ_w satisfying Theorem 2 for each irreducible Coxeter group W of rank five or six. In order to verify Theorem 2, one needs to calculate the characters $\mathrm{Ind}_{C_W(w)}^{W} \varphi_w$ and then confirm that the sum is equal to ρ_S and to $\epsilon\omega_S$. Since these calculations are routine, we display the characters $\mathrm{Ind}_{C_W(w)}^{W}\varphi_w$ (denoted simply as φ_w), ϵ, ρ_S, and ω_S only for the group $W = W(E_6)$.

Because the characters φ_w are one-dimensional, it suffices to list their values on a generating set for $C_W(w)$. For the group $W(E_6)$ generating sets of centralizers are constructed ad hoc. In types B and D we use the notation from [2], which we now briefly review.

The cuspidal classes of $W(B_n)$ are indexed by partitions of n. We always display partitions in non-decreasing order without punctuation. With the labeling of the elements of S given by the diagram  we define the following elements of $W(B_n)$, where we denote the elements of S by $1, 2, \dots, n$ rather than $s_1, s_2, \dots, s_n$ to improve legibility. If $\lambda = \lambda_1\lambda_2\cdots\lambda_k$ is a partition of n then for each $1 \leqslant i \leqslant k$ we define a negative λ_i-cycle

$$c_i = i\,(i-1)\,(i-2)\cdots 212\cdots(i-1)\,i\,(i+1)\cdots(i+\lambda_i)\,.$$

Then each c_i centralizes the element $w_\lambda = c_1c_2\cdots c_k$, which we take to be the representative of the cuspidal class labeled by λ. Whenever $\lambda_i = \lambda_{i+1}$ the element

$$x_i = \prod_{j=0}^{\lambda_i-1}(i+j)\,(i+j-1)\cdots(i+j-\lambda_i)$$

also centralizes w_λ. In fact, if $m(j) = \min\{k \mid \lambda_k = j\}$, then $C_W(w_\lambda)$ is generated by the elements x_i for all i for which $\lambda_i = \lambda_{i+1}$, together with the elements $c_{m(j)}$ for all j appearing as a part of λ. The character φ_{c_λ} of $C_W(w_\lambda)$ is denoted simply by φ_λ.

We view $W(D_n)$ as a reflection subgroup of $W(B_n)$ generated by the reflections $1' = 121$ and $2, 3, \dots, n$. Then $w_\lambda \in W(D_n)$ whenever λ has an even number of parts. In fact, such elements are representatives of the cuspidal classes of $W(D_n)$ and each $C_{W(D_n)}(w_\lambda)$ is the intersection of $W(D_n)$ with $C_{W(B_n)}(w_\lambda)$. The elements c_i are not necessarily elements in $W(D_n)$, but generators of $C_{W(D_n)}(w_\lambda)$ can often be found among products of an even number of c_i.

In each of the following subsections we present the results of our calculations for the finite irreducible Coxeter groups of rank five and six. For each cuspidal class representative w we display a generating set of $C_W(w)$, written as words in the Coxeter generators, and for each generator, the value of the character φ_w. If ζ is an eigenvalue of w on V, we denote the determinant of the representation of $C_W(w)$ on the ζ-eigenspace of w in V by $\det|_\zeta$. If φ_w is a power of $\det|_\zeta$ for some ζ, we also indicate this. By Springer's theory of regular elements [11], the centralizer $C_W(w)$ is a complex reflection group when w is a regular element. When this is the case, we identify $C_W(w)$ as such a group. For $n \geqslant 1$ we denote the n^{th} root of unity $e^{2\pi i/n}$ by ζ_n, the cyclic group of size n by Z_n, and the symmetric group on n letters by S_n.

3.1. $W = W(B_5)$.

λ	Gen	Word	φ_λ	$C_W(w_\lambda)$	Det
$1^3 2$	c_1	1	-1		
	c_4	43212345	-1		
	x_1	2	-1		
	x_2	3	-1		
12^2	c_1	1	-1		
	c_2	2123	-1		
	x_2	4354	-1		
5	c_1	12345	ζ_{10}	Z_{10}	$\det\|_{\zeta_{10}}$
23	c_1	12	-1		
	c_2	3212345	ζ_6		
14	c_1	1	-1		
	c_2	212345	-1		
$1^2 3$	c_1	1	-1		
	c_3	3212345	ζ_6		
	x_1	2	-1		
1^5	S		ϵ	W	$\det\|_{-1}$

3.2. $W = W(\mathsf{B}_6)$.

λ	Gen	Word	φ_λ	$C_W(w_\lambda)$	Det
1^6	S		ϵ	W	$\det\|_{-1}$
$1^4 2$	c_1	1	-1		
	c_5	5432123456	-1		
	x_1	2	-1		
	x_2	3	-1		
	x_3	4	-1		
$1^2 2^2$	c_1	1	-1		
	c_3	321234	-1		
	x_1	2	-1		
	x_3	5465	-1		
2^3	c_1	12	-1	$Z_2 \wr S_3$	
	x_1	3243	-1		
	x_2	5465	-1		
$1^3 3$	c_1	1	-1		
	c_4	432123456	ζ_6		
	x_1	2	-1		
	x_2	3	-1		
123	c_1	1	-1		
	c_2	2123	-1		
	c_3	432123456	ζ_6		
3^2	c_1	123	ζ_6	$Z_2 \wr S_2$	$\det\|_{\zeta_6}$
	x_1	432543654	-1		
$1^2 4$	c_1	1	-1		
	c_2	32123456	-1		
	x_1	2	-1		
24	c_1	12	-1		
	c_2	32123456	-1		
15	c_1	1	-1		
	c_2	2123456	ζ_{10}		
6	c_1	123456	ζ_6	Z_{12}	$(\det\|_{\zeta_{12}})^2$

3.3. $W = W(D_5)$.

λ	Gen	Word	φ_λ	$C_W(w_\lambda)$	Det
$1^3 2$	$w_{1^3 2}$	$1'2321'3431'2345$	ζ_4		
	c_1	$1'$	-1		
	x_1	2	-1		
	x_2	3	-1		
23	w_{23}	$1'3w_{14}$	ζ_{12}		
14	w_{14}	$1'2345$	ζ_8	Z_8	$\det\|_{\zeta_8}$

3.4. $W = W(D_6)$.

λ	Gen	Word	φ_λ	$C_W(w_\lambda)$	Det
$1^2 2^2$	$c_1 c_2$	$1'2$	1		
	$c_1 c_3$	$31'234$	ζ_4		
	x_1	2	-1		
	x_3	5465	-1		
3^2	w_{3^2}	$1343w_{15}$	ζ_3	$G(6,2,2)$	$\det\|_{\zeta_6}$
	c_1^2	$1'323$	ζ_3		
	x_1	432543654	-1		
15	w_{15}	$1'23456$	ζ_5	Z_{10}	$(\det\|_{\zeta_{10}})^2$
$1^3 3$	$c_1 c_2$	$1'2$	1		
	$c_1 c_4$	$43w_{15}$	ζ_3		
	x_1	2	-1		
	x_2	323	-1		
24	w_{24}	$13w_{15}$	ζ_8^3		
	c_2^2	$(321'3456)^2$	ζ_4		
1^6	S		ϵ	W	$\det\|_{-1}$

3.5. $W = W(E_6)$.

Labeling of the elements of S as in the Coxeter graph



,

labeling the conjugacy classes by their Carter diagrams d, and letting r denote the reflection defined by the highest root of W, we define characters $\varphi_d = \varphi_{w_d}$ in the following table.

d	w_d	Gen	φ_d	$C_W(w_d)$	Det
$3A_2$	$12356r$	24542314	ζ_3	G_{25}	$\det\|_{\zeta_3}$
		13	ζ_3		
		56	ζ_3		
$E_6(a_2)$	$w_{E_6}^2$	w_{E_6}	1	G_5	
		234543	ζ_3		
A_5A_1	$13456r$	$w_{A_5A_1}$	ζ_3		
		2345432	-1		
		r	-1		
$E_6(a_1)$	$34w_{E_6}$	$w_{E_6(a_1)}$	ζ_9	Z_9	$\det\|_{\zeta_9}$
E_6	123456	w_{E_6}	-1	Z_{12}	$(\det\|_{\zeta_{12}})^6$

The values of the characters φ_d^W together with ρ_S and ω_S are shown in Table 1.

W	$\emptyset$	$4A_1$	$2A_1$	$3A_2$	A_2	$2A_2$	$D_4(a_1)$	A_3A_1	A_4	$E_6(a_2)$	D_4	A_5A_1	A_22A_1
φ_{3A_2}	80	16	·	−10	−4	2	8	·	·	−2	−2	−2	·
$\varphi_{E_6(a_2)}$	720	16	·	−18	−12	−6	8	·	·	−2	−2	−2	·
$\varphi_{A_5A_1}$	5760	·	·	−72	·	·	·	·	·	·	·	·	·
$\varphi_{E_6(a_1)}$	1440	32	·	−36	12	−3	·	·	·	−4	2	−1	·
φ_{E_6}	4320	96	·	108	·	·	−16	·	·	12	·	·	·
ϵ	1	1	1	1	1	1	1	1	1	1	1	1	1
ρ_S	12320	160	·	−28	−4	−7	·	·	·	4	−2	−5	·
ω_S	12320	160	·	−28	−4	−7	·	·	·	4	−2	−5	·

W	$E_6(a_1)$	E_6	A_1	$3A_1$	A_32A_1	A_3	A_2A_1	$2A_2A_1$	A_5	D_5	A_4A_1	$D_5(a_1)$
φ_{3A_2}	−1	2	·	·	·	·	·	·	·	·	·	·
$\varphi_{E_6(a_2)}$	·	2	·	·	·	·	·	·	·	·	·	·
$\varphi_{A_5A_1}$	·	·	·	·	·	·	·	·	·	·	·	·
$\varphi_{E_6(a_1)}$	·	·	−120	−8	·	·	·	3	1	·	·	·
φ_{E_6}	·	−4	·	·	·	·	·	·	·	·	·	·
ϵ	1	1	−1	−1	−1	−1	−1	−1	−1	−1	−1	−1
ρ_S	−1	·	−120	−8	·	·	·	3	1	·	·	·
ω_S	−1	·	120	8	·	·	·	−3	−1	·	·	·

TABLE 1. The characters φ_d^W, ρ_S, and ω_S for $W = W(E_6)$

4. PROOF OF THEOREM 3

Recall that the normalizer in W of W_L factors as the semidirect product of W_L and a normal complement [8]. When the semidirect product is a direct product,

W_L is called *bulky.* It is shown in [4] that Theorem 3 holds if either W_L is bulky or the rank of W_L is two or less. Also, it is shown in [3] that Theorem 3 holds if W_L is a direct product of Coxeter groups of type A. Thus, to prove Theorem 1 it suffices to prove Theorem 3 for all pairs W, W_L where the rank of W is five or six and L is a proper subset of S for which the following hold.

(1) W_L is not bulky in W,
(2) W_L has rank at least three, and
(3) W_L is not a direct product of Coxeter groups of type A.

After consulting the table of bulky parabolic subgroups in [2], it remains to consider the pairs shown in the next table.

W	W_L			
B_5	A_2B_2			
B_6	A_2B_2,	A_2B_3,	A_3B_2,	$2A_1B_2$
D_5	D_4			
D_6	D_4,	D_5		
E_6	D_4			

In the following subsections, we consider each pair W, W_L separately. For each pair we indicate representatives for the cuspidal conjugacy classes, generators for the centralizers of these representatives, and linear characters of the centralizers that satisfy the conclusion of Theorem 3. For the pairs $W(B_5), W(A_2B_2)$ and $W(E_6), W(D_4)$ we also give the values of $\widetilde{\rho_L}$, $\widetilde{\omega_L}$, α_L, and ϵ. GAP gives the conjugacy classes of a group as a list. The command `ConjugacyClasses` in GAP returns the conjugacy classes of a group as a list. We use the notation w_n to denote the representative of the n^{th} conjugacy class in this list. We denote the longest element of W by w_0 and the longest element in W_L by w_L.

4.1. $W(A_2B_2)$ **in** $W = W(B_5)$**.** Let $L = \{1,2,4,5\}$. The cuspidal conjugacy classes in W_L are represented by w_{13} and w_{15}. The centralizer of w_{15} is $\langle w_{15}\rangle \times \langle w_0\rangle = Z_{12} \times Z_2$. The centralizer of w_{13} is generated by $C_W(w_{15})$ and 1. Define characters $\widetilde{\varphi}_{13}$ of $C_W(w_{13})$ by $(w_{15}, w_0, 1) \mapsto (\zeta_3, 1, -1)$ and $\widetilde{\varphi}_{15}$ of $C_W(w_{15})$ by $(w_{15}, w_0) \mapsto (\zeta_6, 1)$. Then $\widetilde{\varphi}_{13}^{N_W(W_L)} + \widetilde{\varphi}_{15}^{N_W(W_L)}$ equals $\widetilde{\rho_L}$. This character is shown in the following table together with $\widetilde{\omega_L}$, α_L, and ϵ. The conjugacy classes of $N_W(W_L)$ are listed in the order determined by GAP where W is constructed using the command `CoxeterGroup` and $N_W(W_L)$ is constructed using the command `ReflectionSubgroup` and `Normalizer`. The classes are labeled by the orders of their elements and an additional letter to distinguish them from one another.

$N_W(W_L)$	1a	2a	3a	2b	2c	6a	2d	2e	6b	2f	2g	6c	2h	2i	6d
$\widetilde{\rho_L}$	6	·	−3	6	·	−3	−2	·	1	−2	·	1	−2	·	1
$\widetilde{\omega_L}$	6	·	−3	6	·	−3	2	·	−1	2	·	−1	2	·	−1
α_L	1	1	1	−1	−1	−1	1	1	1	−1	−1	−1	1	1	1
ϵ	1	−1	1	−1	1	−1	−1	1	−1	1	−1	1	−1	1	−1

$N_W(W_L)$	2j	2k	6e	4a	4b	12a	4c	4d	12b	2l	2m	6f	2n	2o	6g
$\widetilde{\rho_L}$	−2	·	1	−2	·	1	−2	·	1	6	·	−3	6	·	−3
$\widetilde{\omega_L}$	2	·	−1	−2	·	1	−2	·	1	6	·	−3	6	·	−3
α_L	−1	−1	−1	1	1	1	−1	−1	−1	1	1	1	−1	−1	−1
ϵ	1	−1	1	1	−1	1	−1	1	−1	1	−1	1	−1	1	−1

We see that $\widetilde{\rho_L} = \alpha_L \epsilon \widetilde{\omega_L}$ so that Theorem 3 holds for the pair W, W_L.

4.2. $W(A_2B_2)$ **in** $W = W(B_6)$**.** Let $L = \{1,2,4,5\}$ and let $M = \{1,2,3,4,5\}$. The cuspidal conjugacy classes in W_L are again represented by w_{13} and w_{15}. The centralizer of $w_{15} = 1245$ is $\langle w_{15}\rangle \times \langle w_0\rangle \times \langle w_M\rangle = Z_{12} \times Z_2 \times Z_2$. The centralizer of w_{13} is generated by $C_W(w_{15})$ and 1. Then the characters

$$\widetilde{\varphi}_{13} : (w_{15}, w_0, w_M, 1) \mapsto (\zeta_3, 1, 1, -1)$$
$$\widetilde{\varphi}_{15} : (w_{15}, w_0, w_M) \mapsto (\zeta_6, 1, 1)$$

satisfy Theorem 3 for the pair W, W_L.

4.3. $W(A_2B_3)$ **in** $W = W(B_6)$**.** Let $L = \{1,2,3,5,6\}$. The cuspidal conjugacy classes in W_L are represented by w_{12}, w_{24}, and w_{30}. The centralizer of w_{30} is $\langle 123\rangle \times \langle 56\rangle \times \langle w_0\rangle = Z_6 \times Z_3 \times Z_2$. The centralizer of w_{24} is generated by $w_{24}, 1, w_0$. The centralizer of w_{12} is generated by $C_W(w_{24})$ and $C_W(w_{30})$. The characters

$$\widetilde{\varphi}_{12} : (1, 2, 3, 56, w_0) \mapsto (-1, -1, -1, \zeta_3, -1)$$
$$\widetilde{\varphi}_{24} : (w_{24}, 1, w_0) \mapsto \left(\zeta_3^2, -1, -1\right)$$
$$\widetilde{\varphi}_{30} : (123, 56, w_0) \mapsto \left(-\zeta_3, \zeta_3^2, -1\right)$$

satisfy Theorem 3 for the pair W, W_L.

4.4. $W(A_3B_2)$ **in** $W = W(B_6)$**.** Let $L = \{1,2,4,5,6\}$. The cuspidal conjugacy classes in W_L are represented by w_{23} and w_{25}. The centralizer of w_{25} is $\langle 12\rangle \times \langle 456\rangle \times \langle w_0\rangle = Z_4 \times Z_4 \times Z_2$. The centralizer of w_{25} is generated by $C_W(w_{25})$ and 1. The characters

$$\widetilde{\varphi}_{23} : (1, 2, 456, w_0) \mapsto (-1, -1, \zeta_4, -1)$$
$$\widetilde{\varphi}_{25} : (12, 456, w_0) \mapsto (-1, \zeta_4, -1)$$

satisfy Theorem 3 for the pair W, W_L.

4.5. **$W(2A_1B_2)$ in $W = W(B_6)$.** Let $L = \{1, 2, 4, 6\}$. The cuspidal conjugacy classes in W_L are represented by w_{12} and w_{20}. The centralizer of w_{20} is generated by $12, 4, 6, 5465, r$ where r is the reflection defined by the highest long root of W. The centralizer of w_{12} is generated by $C_W(w_{12})$ and 1. The characters

$$\begin{aligned}\widetilde{\varphi}_{12} &: (1, 2, 4, 6, 5465, r) \mapsto (-1, -1, -1, -1, 1, 1)\\ \widetilde{\varphi}_{20} &: (12, 4, 6, 5465, r) \mapsto (-1, -1, -1, 1, 1)\end{aligned}$$

satisfy Theorem 3 for the pair W, W_L.

4.6. **$W(D_4)$ in $W = W(D_5)$.** Let $L = \{1', 2, 3, 4\}$. The cuspidal conjugacy classes in W_L are represented by w_3, w_9, and w_{11}. (With the notation in the previous section, w_3 is represented by the partition 1111, w_9 by the partition 22, and w_{11} by the partition 13.) The centralizer of w_{11} is $\langle w_{11}\rangle \times \langle w_0\rangle = Z_6 \times Z_2$, the centralizer of w_9 is generated by $x_1 = 232343$ and $1w_0$, and the centralizer of w_3 is generated by W_L and w_0. The characters $\widetilde{\varphi}_3 = \epsilon$ and

$$\begin{aligned}\widetilde{\varphi}_9 &: (x_1, 1w_0) \mapsto (-1, \zeta_4)\\ \widetilde{\varphi}_{11} &: (w_{11}, w_0) \mapsto (\zeta_3, 1)\end{aligned}$$

satisfy Thereom Theorem 3 for the pair W, W_L.

4.7. **$W(D_4)$ in $W = W(D_6)$.** Let $L = \{1', 2, 3, 4\}$ and let $M = \{1', 2, 3, 4, 5\}$. As above, the cuspidal conjugacy classes in W_L are represented by w_3, w_9, and w_{11}. In this case, the centralizer of w_{11} is $\langle w_{11}, 6, w_M\rangle$, the centralizer of w_9 is $\langle 6, x_1, 2w_M\rangle$ where $x_1 = 3243$ is a generator of the centralizer of w_9 in W_L, and the centralizer of w_3 is generated by $C_W(w_9)$ and $C_W(w_{11})$. The characters

$$\begin{aligned}\widetilde{\varphi}_3 &: (1', 2, 3, 4, 6, w_M) \mapsto (-1, -1, -1, -1, 1, 1)\\ \widetilde{\varphi}_9 &: (6, x_1, 2w_M) \mapsto (1, -1, \zeta_4)\\ \widetilde{\varphi}_{11} &: (w_{11}, 6, w_M) \mapsto (\zeta_3, 1, 1)\end{aligned}$$

satisfy Theorem 3 for the pair W, W_L.

4.8. **$W(D_5)$ in $W = W(D_6)$.** Let $L = \{1', 2, 3, 4, 5\}$. The cuspidal conjugacy classes in W_L are represented by w_7, w_{15}, and w_{17}. (With the notation in the previous section, w_7 is represented by the partition 1112, w_{15} by the partition 23, and w_{17} by the partition 14.) The centralizer of w_7 is $\langle w_7, w_0, 1', 2, 3\rangle$, the centralizer of w_{15} is $\langle w_{15}, w_0\rangle$, and the centralizer of w_{17} is $\langle w_{17}, w_0\rangle$. The characters

$$\begin{aligned}\widetilde{\varphi}_7 &: (w_7, w_0, 1', 2, 3) \mapsto (\zeta_4, -1, -1, -1, -1)\\ \widetilde{\varphi}_{15} &: (w_{15}, w_0) \mapsto (\zeta_{12}, -1)\\ \widetilde{\varphi}_{17} &: (w_{17}, w_0) \mapsto (\zeta_8, -1)\end{aligned}$$

satisfy Theorem 3 for the pair W, W_L.

4.9. **$W(D_4)$ in $W = W(E_6)$.** Let $L = \{2, 3, 4, 5\}$. As above, the cuspidal conjugacy classes in W_L are represented by w_3, w_9, and w_{11}. The class containing w_{11} is also labeled by the partition 13. It is convenient to take $y_{13} = 2354$ as a represenative of this class instead of w_{11}. The centralizer of y_{13} is generated by y_{13}, w_M, w_N where $M = \{2, 3, 4, 5, 6\}$ and $N = \{1, 2, 3, 4, 5\}$. (Notice that W_M and W_N are of type D_5 and conjugation by w_M exchanges 2 with 3 and conjugation by w_N exchanges 2 with 5.) The centralizer of w_9 is generated by $x_1 = 4354$, $2w_M$, and $243w_0$. The centralizer of w_3 is generated by $C_W(w_9)$ and $C_W(w_{11})$, and in fact is generated by 3, 4, w_M, and w_0. Define the characters

$$\begin{aligned}\widetilde{\varphi}_3 &: (3, 4, w_M, w_0) \mapsto (-1, -1, 1, 1)\\ \widetilde{\varphi}_9 &: (x_1, 2w_M, 243w_0) \mapsto (-1, -\zeta_4, \zeta_4)\\ \widetilde{\varphi}_{11} &: (y_{13}, w_M, w_N) \mapsto (\zeta_3, 1, 1)\end{aligned}$$

of $C_W(w_3)$, $C_W(w_9)$, and $C_W(w_{11})$ respectively.

In this case, $N_W(W_L) \cong W(F_4)$. Then using the notation from [5, Table C.3] for the irreducible characters of $W(F_4)$ (which is identical to the notation used in CHEVIE), we have

$$\begin{aligned}\widetilde{\varphi}_3^{N_W(W_L)} &= \chi''_{(1,12)}\\ \widetilde{\varphi}_9^{N_W(W_L)} &= \chi'_{(6,6)} + \chi''_{(6,6)}\\ \widetilde{\varphi}_{11}^{N_W(W_L)} &= \chi''_{(2,4)} + \chi_{(9,2)} + \chi''_{(9,6)} + \chi_{(12,4)}\\ \widetilde{\rho_L} &= \chi''_{(1,12)} + \chi''_{(2,4)} + \chi_{(9,2)} + \chi''_{(9,6)} + \chi'_{(6,6)} + \chi''_{(6,6)} + \chi_{(12,4)}\\ \widetilde{\omega_L} &= \chi'_{(1,12)} + \chi'_{(2,16)} + \chi_{(9,10)} + \chi'_{(9,6)} + \chi'_{(6,6)} + \chi''_{(6,6)} + \chi_{(12,4)}.\end{aligned}$$

Now since $\alpha_L \epsilon = \chi_{(1,24)}$ is the sign character, the calculations above show that the characters $\widetilde{\varphi}_3, \widetilde{\varphi}_9, \widetilde{\varphi}_{11}$ above satisfy Theorem 3 for the pair W, W_L.

Acknowledgments: The authors would like to acknowledge support from the DFG-priority program SPP1489 “Algorithmic and Experimental Methods in Algebra, Geometry, and Number Theory”.

References

[1] F. Bergeron, N. Bergeron, R.B. Howlett, and D.E. Taylor. A decomposition of the descent algebra of a finite Coxeter group. *J. Algebraic Combin.*, 1(1): 23–44, 1992.

[2] M. Bishop, J. M. Douglass, G. Pfeiffer, and G. Röhrle. Computations for Coxeter arrangements and Solomon’s descent algebra: Groups of rank three and four. URL `http://arxiv.org/abs/1110.0120`.

[3] J.M. Douglass, G. Pfeiffer, and G. Röhrle. Coxeter arrangements and Solomon’s descent algebra. Preprint, arXiv:1101.2075.

[4] J.M. Douglass, G. Pfeiffer, and G. Röhrle. An inductive approach to Coxeter arrangements and Solomon's descent algebra. *J. Algebraic Combin.*, 35:215–253, 2012.

[5] M. Geck and G. Pfeiffer. *Characters of finite Coxeter groups and Iwahori–Hecke algebras*, volume 21 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 2000.

[6] M. Geck, G. Hiß, F. Lübeck, G. Malle, and G. Pfeiffer. CHEVIE — A system for computing and processing generic character tables. *Appl. Algebra Engrg. Comm. Comput.*, 7:175–210, 1996.

[7] S. Görtzen. Gauss — Extended Gauss Functionality for GAP. 2011. URL http://www.gap-system.org/Packages/gauss.html.

[8] R. B. Howlett. Normalizers of parabolic subgroups of reflection groups. *J. London Math. Soc. (2)*, 21(1):62–80, 1980.

[9] M. Konvalinka, G. Pfeiffer, and C. Röver. A note on element centralizers in finite Coxeter groups. *J. Group Theory*, 14(5):727–745, 2011.

[10] G. Pfeiffer. ZigZag — A GAP3 Package for Descent Algebras of Finite Coxeter Groups. 2011. URL http://schmidt.nuigalway.ie/zigzag.

[11] T. A. Springer. Regular elements of finite reflection groups. *Invent. Math.*, 25:159–198, 1974. ISSN 0020-9910.

Fakultät für Mathematik, Ruhr-Universität Bochum, D-44780 Bochum, Germany
E-mail address: `marcus.bishop@rub.de`

Department of Mathematics, University of North Texas, Denton TX, USA 76203
E-mail address: `douglass@unt.edu`

School of Mathematics, Statistics and Applied Mathematics, National University of Ireland, Galway, University Road, Galway, Ireland
E-mail address: `goetz.pfeiffer@nuigalway.ie`

Fakultät für Mathematik, Ruhr-Universität Bochum, D-44780 Bochum, Germany
E-mail address: `gerhard.roehrle@rub.de`